

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

Armed Services Technical Information Agency

Because of our limited supply, you are requested to return this copy WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters. Your cooperation will be appreciated.

AD

36781

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by
DOCUMENT SERVICE CENTER
KNOTT BUILDING, DAYTON, 2, OHIO

UNCLASSIFIED

AD No. 36 781

ASTIA FILE COPY

STUDIES OF POLYMER ASSOCIATION IN MIXED SOLVENTS

DISTRIBUTION LIST
Technical and Summary Reports

Contract Nonr-839(02)Project NR 330-029

Naval Ordnance Laboratory White Oak Silver Spring, Maryland attn: Dr. Albert Lightbody	(1)	National Advisory Committee for Aeronautics 1724 F Street, N.W. Washington 25, D.C.	(1)
Alleghany Ballistics Laboratory Cumberland, Maryland attn: Dr. Bonner	(1)	Chief, Bureau of Aeronautics Department of the Navy Washington 25, D.C. attn: Code AE-42	(1)
Applied Physics Laboratory The Johns Hopkins University 8261 Georgia Avenue Silver Spring, Maryland attn: Dr. Avery	(1)	Chief, Bureau of Aeronautics Department of the Navy Washington 25, D.C. attn: Code AE-44	(1)
Naval Ordnance Test Station Inyokern China Lake, California attn: Dr. Wiegand attn: Technical Library	(2)	Chief, Bureau of Ordnance Department of the Navy Washington 25, D.C. attn: Code Rela	(1)
Naval Powder Factory Indian Head, Maryland attn: Dr. Thames	(1)	Chief, Bureau of Ordnance Department of the Navy Washington 25, D.C. attn: Code Re2d	(1)
Aerojet Engineering Corp. Azusa, California attn: Dr. Antoni	(1)	Chief, Bureau of Ships Department of the Navy Washington 25, D.C. attn: Code 344	(1)
Reconstruction Finance Corp. Synthetic Rubber Division Research and Development Branch Washington 25, D.C.	(1)	Chief, Bureau of Ships Department of the Navy Washington 25, D.C. attn: Code 346	(1)
Director, National Bureau of Standards Washington 25, D.C. attn: Chief, Organic and Fibrous Materials Division	(1)	Chief, Bureau of Ships Department of the Navy Washington 25, D.C. attn: Code 362	(1)
National Research Council 2101 Constitution Avenue, N.W. Washington 25, D. C. attn: Advisory Board on QM RandD	(1)	Chief, Bureau of Yards and Docks Department of the Navy Washington 25, D.C. attn: Code P300	(1)

DISTRIBUTION LIST (contd)
Technical and Summary Reports

Contract Nonr-839(02)Project NR 330-9

Commanding General Wright Air Development Center Wright-Patterson Air Force Base Dayton, Ohio attn: Materials Laboratory WCRT	(1)	Office of Chief of Ordnance Department of the Army Washington 25, D.C. attn: Code ORDTB	(1)
Commanding General Wright Air Development Center Wright-Patterson Air Force Base Dayton, Ohio attn: Flight Research Laboratory	(1)	Commanding Officer Rock Island Arsenal Rock Island, Illinois attn: R. F. Shaw	(1)
Office of Quartermaster General Department of the Army Washington 25, D.C. attn: Research and Development Div. Chemicals and Plastics Branch	(1)	Commanding Officer Picatinny Arsenal Dover, Delaware attn: ORDTB-T8	(1)
Commanding Officer Signal Corps Engineering Laboratory Department of the Army Fort Monmouth, New Jersey	(1)	The Research and Development Board Pentagon Building Washington 25, D.C. attn: Committee on Materials	(1)
Office of Chief of Engineers Research and Development Division Department of the Army Gravelly Pt., Washington 25, D.C.	(1)	Commander Mare Island Naval Shipyard Rubber Laboratory Vallejo, California	(1)
Engineers Research and Development Laboratory Ft. Belvoir, Virginia attn: Materials Branch, Mr. Mitten	(1)	Dr. J. H. Faull, Jr. 72 Fresh Pond Lane Cambridge 38, Mass.	(1)
		Chief of Naval Research Department of the Navy Washington 25, D.C. attn: Code 424	(2)
		Director Office of Naval Research Branch Office 150 Causeway Street Boston, Mass.	(1)
		Director Office of Naval Research Branch Office 346 Broadway New York 13, N. Y.	(1)

DISTRIBUTION LIST (contd)
Technical and Summary Reports

Contract Nonr-839(02)

Project NR 330-029

Director
Office of Naval Research
Branch Office
844 N. Rush Street
Chicago 11, Illinois (1)

Director
Office of Naval Research
Branch Office
1000 Geary Street
San Francisco 9, Calif. (1)

Director
Office of Naval Research
Branch Office
1030 S. Green Street
Pasadena, California (1)

Officer in Charge
Office of Naval Research
Navy No. 100
Fleet Post Office
New York, New York (2)

Director
Naval Research Laboratory
Washington 25, D.C.
attn: Technical Information
Officer (6)

Armed Services Technical
Information, Document
Service Center,
Knott Building,
Dayton 2, Ohio (5)

Office of Technical Services
Department of Commerce
Washington 25, D.C. (1)

Commanding General
Arnold Engineering
Development Center
Tullahoma, Tenn.
attn: 4507 (1)

TECHNICAL REPORT NUMBER II TO THE NAVY DEPARTMENT
UNDER CONTRACT Nonr839(02) WITH THE OFFICE OF NAVAL
RESEARCH AND THE POLYTECHNIC INSTITUTE OF BROOKLYN

STUDIES OF POLYMER ASSOCIATION IN MIXED SOLVENTS

-- by --

Riad H. Gebren and H. Morawetz
Institute of Polymer Research
Polytechnic Institute of Brooklyn

Project Number NR 330-029
June 21, 1954
Brooklyn 1, New York

STUDIES OF POLYMER ASSOCIATION IN MIXED SOLVENTS

by

Riad H. Gobran and Herbert Morawetz
Institute of Polymer Research, Polytechnic Institute of Brooklyn

Association of small molecules and of polymers.

Molecular association in solution is a fairly common phenomenon, but accurate data about the equilibria have been obtained in comparatively few cases. The dimerization of carboxylic acids in non-polar solvents has probably been studied most extensively, and an exhaustive summary of the results was published recently¹. The association of hydroxylic compounds is somewhat weaker, but the dimerization equilibrium can be studied conveniently by cryoscopic measurements².

While the above associations depend on hydrogen bonding, the association complex between e.g. anthracene and trinitrobenzene is due to the polarization of electrons in the aromatic ring system by the dipoles of the nitro groups³. Of a different type is the widespread occurrence of micelles containing a critical number of ions associated in a single kinetic unit existing in equilibrium with molecularly dispersed ions.

It is not surprising, that the same forces responsible for the association of small molecules produce also association complexes of polymers. This accounts for the dependence of the osmotic or light scattering molecular weights of some polymers on solvent medium and temperature^{4,5,6}. In the case of polyvinyl


chloride, there is undoubtedly a close relationship between its molecular association and the remarkable mechanical properties of the plasticized polymer⁷.

The surprising feature of these polymer associations is the apparent constancy in the size of the molecular aggregates over a wide range of solution concentration. In this respect the polymer aggregates behave like soap micelles, with the "critical micelle concentration" (i.e., the concentration of incipient micelle formation) apparently well below the lowest concentration at which osmotic, or even light scattering measurements are feasible. The formation of micelles has been interpreted convincingly by Debye's theory⁸ as due to the Van der Waals forces, which favor the clustering of the hydrophobic "tails" of the soap ions and the increasing electrostatic work required to add a further unit to a growing micelle. The free energy of the system goes through a sharp minimum for a given number of ions associated in the micelles, and this explains the narrow distribution of their sizes. It is obvious, that this general reasoning is inapplicable, for example, to the association of polystyrene containing a few hydroxyl groups in toluene solution or a methyl methacrylate-methacrylic acid copolymer in benzene.

Spherical aggregates

To illustrate the problem, let us assume that a flexible coil molecule containing a large number of carboxyl groups may be represented by a sphere, with all carboxyls in the interior

of the sphere dimerized by



and only the carboxyls on the surface of the coil in the free state. When the polymers associate, the aggregate is assumed to be again spherical, with the same density as that of the unassociated macromolecule. For D molecules forming an aggregate, the surface area of the complex will be $D^{2/3}$ times the surface of an isolated coil, and the surface area per macromolecule will be changed by a factor $D^{-1/3}$. Thus, if H_0 is the enthalpy of the unassociated polymer molecules, the dependence of enthalpy H on the degree of association D is given by

$$\frac{\partial H}{\partial D} = - \frac{H_0}{3} D^{-4/3} \quad (1)$$

According to the Flory-Huggins theory^{9,10} the entropy of mixing of polymer and solvent is given approximately by

$$\Delta S = -R \{ N_1 \ln \phi_1 + N_2 \ln \phi_2 \} \quad (2)$$

where R is the gas constant, N the number of molecules, ϕ the volume fraction, the indices 1 and 2 referring to solvent and polymer respectively. As the polymer forms aggregates, the volume fractions remain fixed but the number of independent polymer particles decreases, so that

$$\Delta S = R \left\{ N_1 \ln \phi_1 + \frac{N_2}{D} \ln \phi_2 \right\} \quad (3)$$

and the entropy change with degree of association is

$$\partial S / \partial D = RN_2 \ln \phi_2 D^{-2} \quad (4)$$

The rate of change of free energy with degree of association is then given by

$$\frac{\partial F}{\partial D} = \frac{\partial}{\partial D}(H-TS) = -\frac{H_0}{3} D^{-4/3} - RTN_2 \ln \phi_2 D^{-2} \quad (5)$$

For association to occur, $\partial F / \partial D$ must be negative for $D > 1$, or,


$$\frac{H_0}{3} > -RTN_2 \ln \phi_2 \quad (6)$$

It is apparent, that $\partial F / \partial D$ will then be negative for all values of $D > 1$. Thus the model used leads to the conclusion, that once started, polymer association will continue without limit, i.e. to gelation of the system.

"Pearl necklace model" of polymer aggregates.

This conclusion is clearly contrary to experimental evidence, which indicates that although gelation may occur below a critical temperature, above this temperature the polymers associate to complexes of a size determined by temperature and solvent medium, but insensitive to concentration. This behavior can only be explained by assuming, that due to steric factors and intramolecular association, the "effective functionality" of the polymers, i.e., the mean number of groups available for inter-molecular association, is smaller than 2. The model of the association complex can then be represented

schematically as follows:



The two terminal molecules are linked to one neighbor, while all others are linked twice. The mean effective functionality f_0 is related to the number of molecules in the complex D_0 by

$$f_0 = \frac{2+2(D_0-2)}{D_0} = \frac{D_0-1}{D_0} \quad (7)$$

It can easily be verified, that the same result is obtained if the aggregate has a branched structure, provided only that no closed loops are present,

Polymer aggregates in mixed solvents

If a Lewis base is added to the solution of a polymer containing carboxyl groups, a certain fraction of the carboxyls will associate with the base and the number of groups available for inter-polymer association will decrease. This will cause a reduction in the apparent molecular weight of the polymer. The association equilibrium of acid A and base B is given by

$$\frac{(AB)}{(A)(B)} = K \quad (8)$$

and the ratio of the total concentration of acid groups to the concentration of acid groups remaining free in the presence of added base is

$$\frac{(AB) + (A)}{(A)} = 1 + K(B) \quad (9)$$

It is reasonable to postulate, that the effective functionality f of the polymer in a solution containing base

will bear the same relation to the functionality f in the absence of base as the concentration of acid groups unassociated with the base to the total acid group concentration. Utilizing equation (7) and (9)

$$\frac{f_0}{f} = \frac{D_0-1}{D-1} \cdot \frac{D}{D_0} = \frac{(AB)+(A)}{(A)} = 1 + K(B) \quad (10)$$

Thus, the dependence of the degree of association on the concentration of added base should yield information from which the association constant K can be calculated.

Experimental Results

In a previous publication the molecular associations were investigated for mixtures of methyl methacrylate copolymers with methacrylic acid and dimethylaminoethyl methacrylate, respectively¹¹. It was found, that association of the acid copolymer competes strongly with the carboxyl-amine interaction. Also, the importance of any carboxyl association with the ester groups of the copolymers was quite uncertain.

To determine the relative contributions of the different types of association to the formation of the polymer aggregates, the apparent osmotic molecular weight of a methyl methacrylate-methacrylic acid copolymer in benzene solution was measured as a function of added dimethylaminoethyl acetate (DAEA), ethyl acetate, acetic acid and butanone. The copolymer contained 4.9 mole % methacrylic acid (determined by titration with sodium ethoxide in 90 volume % benzene and 10 volume %

ethanol) and its number average molecular weight (measured in pyridine solution, where polymer association cannot take place) was 34,500. The osmotic pressure measurements were carried out as described previously¹¹ at a temperature of 29.8°C. In benzene solution, the degree of association D_0 was found to be 6.86.

The results obtained are listed in Table 1 and plotted in Figure 1. Dimethylaminoethyl acetate and ethyl acetate addition reduced the degree of association of the polymer in the expected manner, and the slopes of the plots of $(D_0-1)D/(D-1)D_0$ against (B) lead to association constants of 86 liters/mole and 6.7 liters/mole respectively. Butanone was relatively inefficient in reducing the degree of polymer association, thirty times as much of it being required to produce the same effect as methyl acetate. While the ester carbonyl would be expected to be more basic than the carbonyl of the ketone, the magnitude of this difference is surprising.

The interpretation of data obtained in solutions to which acetic acid was added is more complex. The acid is largely dimerized, but the monomeric form is chiefly responsible for any interaction with the polymer. The dissociation constant of acetic acid dimer in benzene at 29.8°C. was estimated from published data¹² to be 6.6×10^{-3} moles/liter, and the values of (B) used in Figure 1 refer to the calculated concentrations of the monomeric acid. It can be seen that the plot of $(D_0-1)D/(D-1)D_0$ against (B) is strongly curved in this

case. This is ascribed to the fact, that the monomeric acetic acid associates both with carboxyl and with ester groups of the polymer. In addition, the possibility cannot be excluded that there is some hydrogen bonding of carboxyl with the acetic acid dimer.

Conclusion

It is concluded, that in the acid copolymer used in this and the previous investigation¹¹ most of the aggregation is due to carboxyl-ester association. Also, since the carboxyl-amine association constant is only 13 times as high as the association constant of carboxyl with ester groups, a methacrylate copolymer containing 5 mole % of amine groups would owe somewhat more of its association with a carboxylic polymer to carboxyl-ester than to carboxyl-amine interaction. The method outlined is believed to be a useful tool in determining a variety of association constants, with a possible bearing on the theory of polymer solubility and plasticization¹³.

TABLE 1.

Association of methyl methacrylate-methacrylic acid copolymer in benzene at 29.8°C. in the presence of various additives.


<u>Additive</u>	<u>(B)</u>	<u>D</u>	$\frac{D_0-1}{D-1}$	$\frac{D}{D_0}$
DAEA	0.0062	3.62	1.18	
DAEA	0.0124	1.82	1.89	
DAEA	0.0186	1.52	2.50	
DAEA	0.0310	1.30	3.70	
Methyl acetate	0.012	5.54	1.04	
Methyl acetate	0.021	4.22	1.12	
Methyl acetate	0.042	2.98	1.28	
Butanone	0.553	4.23	1.12	
Acetic Acid (*)	0.0213	5.96	1.03	
Acetic Acid	0.0303	3.84	1.15	
Acetic Acid	0.0375	2.90	1.30	
Acetic Acid	0.0480	1.44	2.80	

(*) Concentrations in monomeric acetic acid.

Bibliography

1. G. Allen and E. F. Caldin, *Quat. Rev.*, 2, 255 (1953).
2. K. L. Wolf, H. Dunken and K. Merkel, *Z. phys. Chem.*, B46, 267 (1940).
3. J. E. Lennard-Jones, *Trans. Far. Soc.*, 32, 1, 37 (1936).
4. P. Doty, H. Wagner and S. Singer, *J. Phys. Colloid Chem.*, 51, 32 (1947).
5. Q. A. Tremontozzi, R. F. Steiner and P. Doty, *J. Am. Chem. Soc.*, 74, 2070 (1952).
6. F. F. Nord, M. Bier and N. Timasheff, *J. Am. Chem. Soc.*, 73, 289 (1951).
7. T. Alfrey, Jr., N. Wiederhorn, R. Stein and A. Tobolsky, *J. Colloid Sci.*, 4, 211 (1949).
8. P. Debye, *Ann. N. Y. Acad. Sci.*, 51, 575 (1949).
9. P. J. Flory, *J. Chem. Phys.*, 10, 51 (1942).
10. M. L. Huggins, *Ann. N. Y. Acad. Sci.*, 43, 1 (1942).
11. H. Morawetz and R. H. Gobran, *J. Polymer Sci.*, 12, 133 (1954).
12. S. Kaufman and C. R. Singleterry, *J. Phys. Chem.*, 56, 604 (1952).
13. A. K. Doolittle, *J. Polymer Sci.*, 2, 121 (1947).

EFFECT OF ADDITIVES ON THE DEGREE OF ASSOCIATION
OF METHYL METHACRYLATE-METHACRYLIC ACID CO-
POLYMER IN BENZENE.



Armed Services Technical Information Agency

Because of our limited supply, you are requested to return this copy WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters. Your cooperation will be appreciated.

AD

36781

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by
DOCUMENT SERVICE CENTER
KNGT BUILDING, DAYTON, 2, OHIO

UNCLASSIFIED